arXiv:1701.03447v1 [math.CO] 10 Jan 2017

A note on near chordal plane triangulations

Sameera M Salam, Nandini J Warrier, Daphna Chacko, K. Murali Krishnan, Sudeep K. S.

Department of Computer Science and Engineering, National Institute of Technology Calicut, Kerala, India 673601

Abstract

It is shown that a plane near triangulated graph is chordal iff it does not contain any induced wheel W_n for $n \geq 5$.

Keywords: Plane triangulated graphs, plane near triangulated graphs, Chordal graphs

1. Introduction

A plane embedding of a (planar) graph G is called a triangulation if the boundary of every face is a cycle of length three. A plane embedding of a simple maximal planar graph is a plane triangulation [1]. If the above definition is relaxed to permit the outer face to have a boundary of length exceeding three, the embedding is called a near triangulation. A graph G is said to contain a chordless cycle if it has an induced cycle C_t for $t \geq 4$. A graph that does not contain any chordless cycle is called a chordal graph [2]. Appolonian networks are well known examples in the literature for maximal planar chordal graphs [3].

In this short note we derive a simple structural characterization for maximal planar chordal graphs that depends only on the neighbourhood of individual vertices. In fact the proof works for near triangulations as well. A proof is presented in Section 3. The following section establishes some notation and definitions.

2. Preliminaries

Let $G = (V, E)$ be a simple undirected graph. A drawing maps each vertex $u \in V$ to a point $\varepsilon(u)$ in $\mathbb{R}^2$ and each edge $(u, v) \in E$ to a path with endpoints $\varepsilon(u)$ and $\varepsilon(v)$. The drawing is a plane embedding if the points are distinct, the paths are simple and do not cross each other and the incidences are limited to the endpoints [2]. The well known Jordan Curve Theorem states that: if J is a simple closed curve in $\mathbb{R}^2$, then $\mathbb{R}^2 - J$ has two components ($Int(J)$ and $Ext(J)$), with J as the boundary of each [4]. Given a plane triangulation G, $Int(G)$ denotes the set of vertices in the interior of the closed curve defined by the boundary of the triangulation.

A wheel W_n of order $n \geq 4$ is the graph $K_1 \cup C_{n-1}$. The cycle C_{n-1} is called the rim of the wheel [2]. A separator in a connected graph is a set of vertices, the removal of which disconnects

Email addresses: shemi.nazir@gmail.com (Sameera M Salam), nandini.wj@gmail.com (Nandini J Warrier), daphna.chacko@gmail.com (Daphna Chacko), kmurali@nitc.ac.in (K. Murali Krishnan), sudeep@nitc.ac.in (Sudeep K. S.)

the graph. A clique in a graph is a set of pairwise adjacent vertices. A clique separator is a separator which is a clique [5].

Definition 2.1 (Separating triangle)**.** *A clique separator with three vertices is called a separating triangle.*

Definition 2.2 ($W-$graph)**.** *A near triangulation G, $G \neq K_4$ is called a $W-$graph if G does not contain a separating triangle.*

Let G be any graph and $v \in V(G)$. Define the open neighbourhood of the vertex v as $N(v)$= $\{v' \in V : (v', v) \in E\}$. The closed neighbourhood of v, $N[v] = \{v\} \cup N(v)$.

3. Non chordal plane near triangulated graphs

We first prove the following Lemma:

Lemma 3.1. *If G is a $W-$graph then $\forall u \in Int(G)$, $N[u]$ induces a wheel W_k for some $k \geq 5$.*

Proof. Let G be a $W-$graph, $u \in Int(G)$. As G is plane triangulated, it is easy to see that $|N(u)| \geq 3$. Moreover as G is a $W-$graph and $G \neq K_4$, $|N(u)| \neq 3$ for otherwise vertices in $N(u)$ will form a separating triangle. Hence let $N(u) = \{u_1, u_2, \ldots u_{k-1}\}$ for some $k \geq 5$ and let G' be the subgraph of G induced by $N(u)$. To complete the proof it is enough to prove that G' induces a cycle in G. We rule out the following three cases.

Case 1 : Suppose G' is not connected and has components $G_1, G_2, \ldots G_l$ where $2 \leq l \leq k-1$. Without loss of generality we may assume that G_1 contains vertices $\{u_1, u_2, \ldots u_i\}$ and G_2 contains vertices $\{u_{i+1}, \ldots u_j\}$ $(1 \leq i < j \leq k-1)$ in the clockwise order. Then as G is triangulated with $u \in Int(G)$ and $\{(u, u_i), (u, u_{i+1})\} \subseteq E(G)$, it must be the case that $(u_i, u_{i+1}) \in E(G)$, for otherwise u, u_i and u_{i+1} will be part of a face of length exceeding three, contradicting the assumption that u_i and u_{i+1} are in different components.

Case 2 : Suppose G' is a path. With out loss of generality we may assume that the vertices in the path are $u_1, u_2, \ldots . u_{k-1}$ ordered from left to right. As $\{(u, u_1), (u, u_{k-1})\} \subseteq E(G)$ and $u \in Int(G)$, it follows that G must have an internal face f containing edges u, u_1 and u_{k-1}. But as $(u_1, u_{k-1}) \notin E(G)$, f cannot be a triangle, a contradiction.

Case 3 : Suppose G' contains a vertex u_i with degree at least three. Let x, y and z be any three neighbours of u_i in G'. Then one among the triplets (u, u_i, x), (u, u_i, y) and (u, u_i, z) will be a separating triangle in G, a contradiction.

It follows therefore that G' induces a cycle of length $k-1$

□

The observation below follows directly from the definition of a separating triangle.

Observation 3.1. *Let the vertices u, v and w form a separating triangle in a plane near triangulated graph G. Then the triangle (u, v, w) separates G into two components (say G_1 and G_2) with (u, v, w) as the boundary. Moreover, G is chordal iff both G_1 and G_2 are chordal.*

The following theorem shows that chordal plane near triangulations are characterized by the closed neighbourhood of internal vertices.

Theorem 3.1. *A plane near triangulated graph is not chordal iff it contains an induced wheel W_k for some $k \geq 5$.*

Proof. Let $G(V, E)$ be a near triangulated graph. If G contains an induced W_k for some $k \geq 5$ then the rim of W_k is a chordless cycle and G is non chordal.

Conversely, suppose G is non chordal. If G is a $W-$graph, then the claim follows from Lemma 3.1. Otherwise, let $\triangle = (u, v, w)$ be a separating triangle in G. Then by the Observation 3.1, (u, v, w) divides G into two components say G_1 and G_2 and either G_1 or G_2 or both are not chordal. The claim follows by induction. □

Acknowledgment

We would like to thank Prof Ajith Diwan, IIT Bombay for his valuable suggestions and support.

References

[1] R. Balakrishnan, K. Ranganathan, A textbook of graph theory, Springer Science & Business Media, 2012.

[2] D. B. West, et al., Introduction to graph theory, Vol. 2, Prentice hall Upper Saddle River, 2001.

[3] P. S. Kumar, C. E. V. Madhavan, A new class of separators and planarity of chordal graphs, in: Proceedings of the Ninth Conference on Foundations of Software Technology and Theoretical Computer Science, Springer-Verlag, London, UK, UK, 1989, pp. 30–43.
URL `http://dl.acm.org/citation.cfm?id=646827.707381`

[4] W. Fulton, Algebraic topology: a first course, Vol. 153, Springer, 1995.

[5] A. Berry, R. Pogorelcnik, G. Simonet, An introduction to clique minimal separator decomposition, Algorithms 3 (2) (2010) 197–215.